Plant Pathology

INDEX
VOLUME 30, 1981

Author: *pages* iii–iv
Subject: *pages* v– ix

AUTHOR INDEX

Alford, D. V. *(joint author)*, see Graham, C. W., 141–5.

Bartlett, P. W. Colorado beetles reported in England, Wales and Scotland, 1980, 41–6.

Bartlett, P. W. *Trionymus diminutus* (Leonardi) (Homoptera: Pseudococcidae) infesting New Zealand flax in England, 56–8.

Bartlett, P. W. and Powell, D. F. Introduction of American serpentine leaf miner, *Liriomyza trifolii*, into England and Wales and its eradication from commercial nurseries, 1977–81, 185–93.

Boesewinkel, H. J. Two species of powdery mildew on exotic rhododendrons in Edinburgh, 119–20.

Brain, P. J. and Whittington, W. J. The influence of soil pH on the severity of swede powdery mildew infection, 105–9.

Brown, D. J. F. *(joint author)*, see Thomas, P. R., 147–51.

Buckler, Eva *(joint author)*, see Sellwood, Jennifer E., 179–80.

Burges, H. D., *(joint author)*, see Kanagaratnam, P., 117–18.

Carter, Margaret A. *(joint author)*, see Locke, T., 89–93.

Chambers, D. A. *(joint author)*, see Edney, K. L., 167–70.

Channon, A. G. and Thomson, Moira C. Parsnip canker caused by *Cylindrocarpon destructans*, 181.

Clarkson, J. D. S. Relationship between eyespot severity and yield loss in winter wheat, 125–31.

Clayden, Ingrid and Hooper, D. J. New weed hosts for the giant race of *Ditylenchus dipsaci* (Kühn) Filipjev, 251–2.

Connolly, G. P. and Fletcher, J. T. Procedures for the generation of formaldehyde vapour to fumigate structures, 245–50.

Cooper, D. A. Forecasting egg populations of wheat bulb fly, *Delia coarctata* (Fall.), from catches of adult females in water-traps, 31–5.

Cooper, P. Strawberry latent ringspot virus in parsnips, 59–60.

Cotten, J. and Roberts, Helen. Root-lesion nematodes (*Pratylenchus* spp.) in raspberry-cane beds entered for certification in England, 101–4.

Dennis, E. B. *(joint author)*, see Green, C. D., 65–71.

Edney, K. L. and Chambers, D. A. The use of metalaxyl to control *Phytophthora syringae* rot of apple fruits, 167–70.

Evers, A. D. *(joint author)*, see King, J. E., 51–3.

Ewart, J. M. *(joint author)*, see Sellwood, Jennifer E., 179–80.

Fletcher, J. T. *(joint author)*, see Connolly, G. P., 245–50.

Gay, Caroline E. *(joint author)*, see Locke, T., 89–93.

Gibson, R. W. Rapid spread by mites of ryegrass mosaic virus from old sward to seedling ryegrass and its prevention by aldicarb, 25–9.

Gibson, R. W. and Gunenc, Yildiz. Effects of covering potato crops with clear polyethylene film on spread of potato virus Y, 233–5.

Gladders, P. Purple blotch of leeks caused by *Alternaria porri*, 61.

Gould, H. J. and Jessop, N. Field tests with acaricides for the control of *Tetranychus urticae* Koch on strawberries, 171–5.

Graham, C. W. and Alford, D. V. The distribution and importance of cabbage stem flea beetle (*Psylliodes chrysocephala* (L.)) on winter oilseed rape in England, 141–5.

Green, C. D. and Dennis, E. B. An analysis of the variability in yield of pea crops attacked by *Heterodera goettingiana*, *Helicotylenchus vulgaris* and *Pratylenchus thornei*, 65–71.

Greenway, A. R. *(joint author)*, see Wall, C., 73–6.

Gunenc, Yildiz *(joint author)*, see Gibson, R. W., 233–5.

Hall, R. A. *(joint author)*, see Kanagaratnam, P., 117–18.

Halstead, A. J. A whitefly pest of sweet bay, 123.

Hammond, John. Viruses occurring in *Plantago* species in England, 237–43.

Harris, D. C. and Stickels, J. E. Crown rot (*Phytophthora cactorum*) in glasshouse-grown strawberries at East Malling Research Station, 205–12.

Harrison, J. G. Chocolate spot of field beans in Scotland, 111–15.

Harrison, J. G. *Fusarium acuminatum* on field bean, 121.

Hooper, D. J. *(joint author)*, see Clayden, Ingrid, 251–2.

Humphreys-Jones, D. R. Stem rot (*Microdiplodia ceanothi*) of *Ceanothus dentatus*, 62–3.

Humphreys-Jones, D. R. *Coniothyrium lavandulae* wilt and die back of *Lavandula angustifolia*, 183.

Jacobson, R. J. and Thomas, K. P. *Rhyzopertha dominica* (F.) (Coleoptera: Bostrichidae) in stored home-grown barley, 54–5.

Jessop, N. *(joint author)*, see Gould, H. J., 171–5.

Jones, F. G. W. *(joint author)*, see Webley, D. P., 217–24.

Jones, I. T. *(joint author)*, see Potter, L. R., 133–9.

Jones, R. A. C. *(joint author)*, see Torrance, L., 1–24.

Jones, R. A. C. Use of continuous light to improve diagnosis of potato leaf roll virus in tuber indexing tests, 225–32.

Jordan, V. W. L. Aetiology of barley net blotch caused by *Pyrenophora teres* and some effects on yield, 77–87.

Kanagaratnam, P., Hall, R. A. and Burges, H. D. Effect of fungi on the black currant gall mite, *Cecidophyopsis ribis*, 117–18.

King, J. E., Evers, A. D. and Stewart, B. A. Black-point of grain in spring wheats of the 1978 harvest, 51–3.

Locke, T., Thorpe, I. G., Carter, Margaret A. and Gay, Caroline E. Differential sensitivity of *Pyrenophora teres* to organomercury formulations, 89–93.

Port, C. M. and Scopes, N. E. A. Biological control by predatory mites (*Phytoseiulus persimilis* Athias-Henriot) of red spider mite (*Tetranychus urticae* Koch) infesting strawberries grown in 'walk-in' plastic tunnels, 95–9.

Potter, L. R. and Jones I. T. Interaction between barley yellow dwarf virus and powdery mildew in four barley genotypes, 133–9.

Powell, D. F. Fumigation of soil with methyl bromide to kill diapausing Colorado beetles, 159–65.

Powell, D. F. *(joint author)*, see Bartlett, P. W., 185–93.

Powell, D. F. The eradication campaign against American serpentine leaf miner, *Liriomyza trifolii*, at Efford Experimental Horticulture Station, 195–204.

Roberts, Helen *(joint author)*, see Cotten, J., 101–4.

Roberts, Helen. New or unusual host-plant records for plant-parasitic nematodes, 1977–80, 182.

Savage, M. J. *(joint author)*, see Williams, J. J. W., 122.

Scopes, N. E. A. *(joint author)*, see Port, C. M., 95–9.

Sellwood, Jennifer E. Vascular blackening of chicory caused by a pectolytic isolate of *Pseudomonas fluorescens*, 179–80.

Smallshire, D. *(joint author)*, see Williams, J. J. W., 122.

Stewart, B. A. *(joint author)*, see King, J. E., 51–3.

Stickels, J. E. *(joint author)*, see Harris, D. C., 205–12.

Stroyan, H. L. G. A North American lupin aphid found in Britain, 253.

Thomas, K. P. *(joint author)*, see Jacobson, R. J., 54–5.

Thomas, P. R. and Brown, D. J. F. Some hosts of a *Meloidogyne ardenensis* population found in Scotland, 147–51.

Thomson, Moira C. *(joint author)*, see Channon, A. G., 181.

Thorpe, I. G. *(joint author)*, see Locke, T., 89–93.

Torrance, L. and Jones, R. A. C. Recent developments in serological methods suited for use in routine testing for plant viruses, 1–24.

Torrance, L. Use of forced buds to extend the period of serological testing in surveys for fruit tree viruses, 213–16.

Wall, C. and Greenway, A. R. An effective lure for use in pheromone traps for monitoring pea moth, *Cydia nigricana* (F.), 73–6.

Webley, D. P. and Jones, F. G. W. Observations on *Globodera pallida* and *G. rostochiensis* on early potatoes, 217–24.

White, P. F. Control of the mushroom phorid *Megaselia halterata* (Wood) with dichlorvos applied by an automatic atomiser, 37–40.

Whittington, W. J. *(joint author)*, see Brain, P. J., 105–9.

Williams, J. J. W., Savage, M. J. and Smallshire, D. *Longidorus vineacola* Sturhan & Weischer damaging spring barley and onions in England, 122.

Wooster, P. Post-harvest tuber indexing in the seed potato certification schemes of England and Wales, 1978, 153–8.

SUBJECT INDEX

Abacarus hystrix, 25.
Acremoniella, 51.
Agromyza albipennis, 199.
Agromyza felleri, 199.
Agromyza nigrella, 199.
Agromyza spp., 199.
Aldicarb:
  for Control of American serpentine leaf miner, 188, 198.
  for Control of red spider mite on strawberry, 171.
  for Control of *Trionymus diminutus* on New Zealand flax, 56.
  for Prevention of spread of ryegrass mosaic virus, 25.
Alternaria porri, 61.
Alternaria spp., 51.
Amitraz:
  for Control of red spider mite on strawberry, 171.
Anemone:
  Meloidogyne ardenensis on poppy anemone, 147.
Aphelenchoides fragariae, 182.
Aphelenchoides ritzemabosi, 182.
Aphis nasturtii, 234.
Apple:
  Use of forced buds to extend period of serological testing in surveys for fruit tree viruses, 213.
  Use of metalaxyl to control *Phytophthora syringae* rot of fruits, 167.
Ascochyta, 51.
Ash:
  Host of *Meloidogyne ardenensis*, 147.
Aspergillus, 51.
Aspergillus fumigatus, 55.
Aureobasidium spp., 51.

Barley:
  Aetiology of net blotch caused by *Pyrenophora teres* and some effects on yield, 77.
  Differential sensitivity of *Pyrenophora teres* to organomercury formulations, 89.
  Host of *Meloidogyne ardenensis*, 147.
  Interaction between barley yellow dwarf virus and powdery mildew in four barley genotypes, 133.
  Longidorus vineacola damaging spring barley in England, 122.
  Rhyzopertha dominica in stored home-grown barley, 54.
Bay:
  Whitefly pest of sweet bay, 123.
Beans:
  American serpentine leaf miner on navy bean and runner bean, 200.
  Chocolate spot of field beans in Scotland, 111.
  Fusarium acuminatum on field bean, 121.
Beetroot:
  Host of *Meloidogyne ardenensis*, 147.
Bemisia hancocki, 123.
Benomyl:
  Effect on control of *Phytophthora syringae* of apple fruits with metalaxyl, 167.
Biological Control:
  Red spider mite infesting strawberries grown in 'walk-in' plastic tunnels, 95.
Botrytis, 52.
Botrytis cinerea, 51, 113, 245.
Botrytis fabae, 111.

Carrot:
  Host of *Meloidogyne ardenensis*, 147.
Ceanothus:
  Stem rot of *Ceanothus dentatus*, 62.
Cecidophyopsis ribis, 117.
Cherry:
  Use of forced buds to extend period of serological testing in surveys for fruit tree viruses, 213.
Chicory:
  Vascular blackening caused by a pectolytic isolate of *Pseudomonas fluorescens*, 179.
Chlorpyrifos:
  for Control of American serpentine leaf miner, 201.
  for Control of red spider mite on strawberry, 171.
Chrysanthemum:
  Eradication campaign against American serpentine leaf miner at Efford Experimental Horticulture Station, 195.
  Introduction of American serpentine leaf miner into England and Wales and its eradication from commercial nurseries, 1977–81, 185.
Cladosporium herbarum, 183.
Cladosporium spp., 51, 113.
Colorado Beetle:
  Beetles reported in England, Wales and Scotland, 1980, 41.
  Fumigation of soil with methyl bromide to kill diapausing beetles, 159.
Coniothyrium lavandulae, 183.
Cornflower:
  Host of American serpentine leaf miner, 200.
Cucumber:
  Host of *Meloidogyne ardenensis*, 147.
Cydia nigricana, 73.
Cyhexatin:
  for Control of red spider mite on strawberry, 171.
Cylindrocarpon destructans, 181.

Dahlia:
Host of American serpentine leaf miner, 200.
DDT:
for Control of American serpentine leaf miner, 201.
Delia coarctata, 31.
Demeton-S-methyl:
for Control of red spider mite on strawberry, 171.
for Control of *Trionymus diminutus* on New Zealand flax, 56.
Diazinon:
for Control of American serpentine leaf miner, 188.
for Control of mushroom phorid, 37.
for Control of *Trionymus diminutus* on New Zealand flax, 56.
Dichlorvos:
for Control of mushroom phorid, 37.
Dicofol:
for Control of red spider mite on strawberry, 171.
Dienochlor:
for Control of red spider mite on strawberry, 171.
Dimethoate:
for Control of red spider mite on strawberry, 171.
for Control of *Trionymus diminutus* on New Zealand flax, 56.
Dinobuton:
for Control of red spider mite on strawberry, 171.
Dioxathion:
for Control of red spider mite on strawberry, 171.
Disease Assessment:
Chocolate spot of field beans, 112.
Eyespot in winter wheat, 125.
Potato virus Y and potato leaf roll virus infections, 153.
Potato virus Y infection on potato tubers, 234.
Swede powdery mildew infection, 105.
Disorder:
Black-point of grain in spring wheats of 1978 harvest, 51.
Ditylenchus dipsaci, 182, 251.

Elder:
Host of *Meloidogyne ardenensis*, 147.
Epicoccum, 52.
Erysiphe cruciferarum, 105, 119.
Erysiphe graminis, 133.

Fenbutatin oxide:
for Control of red spider mite on strawberry, 171.
Fenitrothion:
for Control of *Rhyzopertha dominica* in stored barley, 54.
for Control of American serpentine leaf miner, 188.
Flax:
Trionymus diminutus infesting New Zealand flax in England, 56.
Forecasting:
Egg populations of wheat bulb fly from catches of adult females in water-traps, 31.
Formaldehyde:
Procedures for generation of vapour to fumigate structures, 245.
Formetanate:
for Control of red spider mite on strawberry, 171.
Fusarium, 52.
Fusarium acuminatum, 121.
Fusarium nivale, 51.

Gamma-HCH:
for Control of American serpentine leaf miner, 188, 198, 201.
Gerbera:
Introduction of American serpentine leaf miner into England and Wales and its eradication from commercial nurseries, 1977–81, 185.
Globodera pallida, 217.
Globodera rostochiensis, 217.
Grass:
Meloidogyne ardenensis on annual meadow-grass, 147.

Hawthorn:
Host of American serpentine leaf miner, 200.
Helicotylenchus vulgaris, 65.
Helminthosporium, 51.
Heptenophos:
for Control of American serpentine leaf miner, 188, 198, 201.
Heterodera goettingiana, 65.
Hirsutella thompsonii, 117.

Iprodione:
for Control of purple blotch of leeks, 61.
Itersonilia pastinacae, 181.

Lavandula:
Wilt and die back of *Lavandula angustifolia* caused by *Coniothyrium lavandulae*, 183.
Leek:
Purple blotch caused by *Alternaria porri*, 61.
Leptinotarsa decemlineata, 41, 159.
Liriomyza eupatorii, 199.
Liriomyza flaveola, 199.
Liriomyza lutea, 199.
Liriomyza orbona, 199.
Liriomyza pusilla, 199.
Liriomyza spp., 199.
Liriomyza strigata, 199.

Liriomyza trifolii, 185, 195.
Longidorus vineacola, 122.
Losses caused by:
Barley net blotch, 85.
Eyespot in winter wheat, 125.
Heterodera goettingiana, *Helicotylenchus vulgaris* and *Pratylenchus thornei* in pea, 65.
Lupin:
North American lupin aphid found in Britain, 253.
Lyonetia clerckella, 199.

Macrosiphum albifrons, 253.
Macrosiphum euphorbiae, 233.
Malathion:
for Control of *Trionymus diminutus* on New Zealand flax, 57.
Mancozeb:
for Control of purple blotch of leeks, 61.
Marigold:
American serpentine leaf miner on French marigold and pot marigold, 200.
Megaselia halterata, 37.
Meloidogyne ardenensis, 147.
Metalaxyl:
for Control of *Phytophthora syringae* rot of apple fruits, 167.
Metarhizium anisopliae, 118.
Methiocarb:
for Control of red spider mite on strawberry, 171.
Methyl bromide:
for Control of American serpentine leaf miner, 188, 198.
to Kill diapausing Colorado beetles, 159.
Microdiplodia ceanothi, 62.
Mushroom:
Control of mushroom phorid with dichlorvos applied by automatic atomiser, 37.
Myzus persicae, 233.

Nasturtium:
Host of American serpentine leaf miner, 200.
Nematodes:
Analysis of variability in yield of pea crops attacked by *Heterodera goettingiana*, *Helicotylenchus vulgaris* and *Pratylenchus thornei*, 65.
Longidorus vineacola damaging spring barley and onions in England, 122.
New or unusual host-plant records for plant-parasitic nematodes, 182.
New weed hosts for giant race of *Ditylenchus dipsaci*, 251.
Observations on *Globodera pallida* and *G. rostochiensis* on early potatoes, 217.
Root-lesion nematodes in raspberry-cane beds entered for certification in England, 101.
Some hosts of a *Meloidogyne ardenensis* population in Scotland, 147.
Nepticula aurella, 199.
Nepticula sp., 199.
Nicotine:
for Control of American serpentine leaf miner, 188.

Oat:
Host of *Meloidogyne ardenensis*, 147.
Oilseed Rape:
Distribution and importance of cabbage stem flea beetle on winter oilseed rape in England, 141.
Omethoate:
for Control of red spider mite on strawberry, 171.
Onion:
Host of *Meloidogyne ardenensis*, 147.
Longidorus vineacola damaging onions in England, 122.
Organomercury compounds:
Differential sensitivity of *Pyrenophora teres* to organomercury formulations, 89.
Oxamyl:
for Control of American serpentine leaf miner, 188, 198.
for Control of nematodes in pea, 66.

Parsnip:
Canker caused by *Cylindrocarpon destructans*, 181.
Host of American serpentine leaf miner, 200.
Strawberry latent ringspot virus, 59.
Pea:
Analysis of variability in yield of crops attacked by *Heterodera goettingiana*, *Helicotylenchus vulgaris* and *Pratylenchus thornei*, 65.
Host of American serpentine leaf miner, 200.
Penicillium, 52.
Permethrin:
for Control of American serpentine leaf miner, 188, 198.
Pest Assessment:
Aphids on potato foliage, 233.
Phenisobromolate:
for Control of red spider mite on strawberry, 173.
Phorate:
for Control of aphids on potato, 233.
for Control of potato virus Y, 233.
Phytomyza horticola, 199.
Phytomyza nigra, 199.
Phytomyza pastinacae, 199.
Phytomyza ranunculi, 199.
Phytomyza spp., 199.

Phytomyza syngenesiae, 186, 199.
Phytophthora cactorum, 205.
Phytophthora syringae, 167.
Phytoseiulus persimilis, 95.
Pirimiphos-methyl:
for Control of red spider mite on strawberry, 171.
Plantago:
Viruses occurring in *Plantago* species in England, 237.
Plum:
Use of forced buds to extend period of serological testing in surveys for fruit tree viruses, 213.
Potato:
Effects of covering crops with clear polyethylene film on spread of potato virus Y, 233.
Host of American serpentine leaf miner, 200.
Observations on *Globodera pallida* and *G. rostochiensis* on early potatoes, 217.
Post-harvest tuber indexing in seed potato certification schemes of England and Wales, 153.
Use of continuous light to improve diagnosis of potato leaf roll virus in tuber indexing tests, 225.
Pratylenchus crenatus, 101.
Pratylenchus fallax, 101.
Pratylenchus neglectus, 101.
Pratylenchus penetrans, 101, 182.
Pratylenchus thornei, 65, 101.
Predators:
Phytoseiulus persimilis, 95.
Propargite:
for Control of red spider mite on strawberry, 171.
Propiconazole:
for Control of barley net blotch, 77.
Pseudocercosporella herpotrichoides, 125.
Pseudomonas fluorescens, 179.
Psylliodes chrysocephala, 141.
Puccinia horiana, 185.
Pynosect:
for Control of mushroom phorid, 38.
Pyrenophora teres, 77, 89.
Pyrethrum:
for Control of whitefly on sweet bay, 123.

Quinalphos:
for Control of red spider mite on strawberry, 171.
Quinomethionate:
for Control of red spider mite on strawberry, 171.

Raspberry:
Host of *Meloidogyne ardenensis*, 147.
Root-lesion nematodes in cane beds entered for certification in England, 101.
Resistance to Pesticides:
Red spider mite to dicofol and organophosphorus pesticides, 171.
Rhyzopertha dominica to gamma-HCH, 55.
Resmethrin:
for Control of American serpentine leaf miner, 188, 198.
for Control of whitefly on sweet bay, 123.
Rhododendron:
Two species of powdery mildew on exotic rhododendron, 119.
Rhynchaenus fagi, 199.
Rhyzopertha dominica, 54.
Rose:
Meloidogyne ardenensis on dog rose, 147.
Ryegrass:
Meloidogyne ardenensis on Italian ryegrass, 147.
Rapid spread by mites of ryegrass mosaic virus from old sward to seedling ryegrass and its prevention by aldicarb, 25.

Septoria, 52.
Sloe:
Host of *Meloidogyne ardenensis*, 147.
Snapdragon:
Host of American serpentine leaf miner, 200.
Sphaerotheca pannosa, 119.
Steam:
for Control of American serpentine leaf miner, 188, 198.
Stemphylium, 51.
Stemphylium botryosum, 61.
Stemphyllium sp., 183.
Strawberry:
Biological control by predatory mites of red spider mite in 'walk-in' plastic tunnels, 95.
Crown rot in glasshouse-grown strawberries at East Malling Research Station, 205.
Field tests with acaricides for control of *Tetranychus urticae*, 171.
Host of *Meloidogyne ardenensis*, 147.
Surveys:
Cabbage stem flea beetle on winter oilseed rape in England, 141.
Fruit tree viruses, 213.
Fungal pathogens of black currant gall mite in England, 117.
Host plants of American serpentine leaf miner, 195.
Infestations of American serpentine leaf miner, 185, 195.
Viruses occurring in *Plantago* species in England, 237.
Swede:
Influence of soil pH on severity of powdery mildew infection, 105.
Sweet pea:
Host of American serpentine leaf miner, 200.
Sycamore:
Host of *Meloidogyne ardenensis*, 147.

Tetradifon:
for Control of red spider mite on strawberry, 173.
Tetranychus urticae, 95, 171.
Thiofanox:
for Control of red spider mite on strawberry, 171.
Tomato:
Host of American serpentine leaf miner, 200.
Host of *Meloidogyne ardenensis*, 147.
Traps and Trapping:
Effective lure for use in pheromone traps for monitoring pea moth, 73.
Forecasting egg populations of wheat bulb fly from catches of adult females in water-traps, 31.
Monitoring adult American serpentine leaf miner with yellow sticky traps, 196.
Triazophos:
for Control of American serpentine leaf miner, 188.
for Control of red spider mite on strawberry, 171.
Trichlorphon:
for Control of American serpentine leaf miner, 201.
for Control of *Trionymus diminutus* on New Zealand flax, 57.
Trichothecium roseum, 51.
Trionymus diminutus, 56.
Tropaeolum peregrinum:
Host of American serpentine leaf miner, 200.

Valerian:
American serpentine leaf miner on red valerian, 200.
Verticillium lecanii, 117.
Viruses:
Effects of covering potato crops with clear polyethylene film on spread of potato virus Y, 233.
Interaction between barley yellow dwarf virus and powdery mildew in four barley genotypes, 133.
Occurring in *Plantago* species in England, 237.
Post-harvest tuber indexing in seed potato certification schemes of England and Wales, 153.
Rapid spread by mites of ryegrass mosaic virus from old sward to seedling ryegrass and its prevention by aldicarb, 25.
Recent developments in serological methods suited for use in routine testing for plant viruses, 1.
Strawberry latent ringspot virus in parsnips, 59.
Use of continuous light to improve diagnosis of potato leaf roll virus in tuber indexing tests, 225.
Use of forced buds to extend period of serological testing in surveys for fruit tree viruses, 213.

Weather in England and Wales:
Spring and Summer 1980, 49.
Autumn and Winter, 1980–81, 177.
Wheat:
Black-point of grain in spring wheats of 1978 harvest, 51.
Host of *Meloidogyne ardenensis*, 147.
Relationship between eyespot severity and yield loss in winter wheat, 125.

Zineb:
for Control of purple blotch of leeks, 61.